

Parametrization of the QCD coupling in Hard and Regge processes

B.I. Ermolaev

Ioffe Physico-Technical Institute, 194021 St. Petersburg, Russia

S.I. Troyan

St. Petersburg Institute of Nuclear Physics, 188300 Gatchina, Russia

We examine the parametrization of the QCD coupling in the Bethe-Salpeter equations for the hard and Regge processes and determine the argument of α_s of the factorized gluon. Our analysis shows that for the hard processes $\alpha_s = \alpha_s(k_\perp^2/(1-\beta))$ where $k_\perp^2$ and β are the longitudinal and transverse moment of the soft parton. On the other hand, in the Regge processes $\alpha_s = \alpha_s(k_\perp^2/\beta)$. We have also shown that the well-known parametrization $\alpha_s = \alpha_s(k_\perp^2)$ in the DGLAP equations stands only if the lowest integration limit μ^2 over $k_\perp^2$ (the starting point of the Q^2 -evolution) obeys the relation $\mu \gg \Lambda_{QCD} \exp(\pi/2)$, otherwise the coupling should be replaced by the more complicated expression.

PACS numbers: 12.38.Cy

I. INTRODUCTION

Total resummations of radiative corrections in QCD are often performed with composing and solving evolution equations. Quite often such equations are of the Bethe-Salpeter type, with one gluon (or one ladder rung) being factorized. In particular, both the DGLAP and BFKL equations are of that type. As is well-known, the argument of α_s in the DGLAP equations is Q^2 . It follows from the dependence

$$\alpha_s = \alpha_s(k_\perp^2) \quad (1)$$

in each rung of the ladder Feynman graphs. Such a dependence originally was used in analogy to the parametrization of α_s in the hard kinematics according to the results of Refs. [1, 2]. Later, the proof of the parametrization (1) in the DGLAP context was suggested in Ref. [3]. However, that proof was not done accurate enough. In our recent paper Ref. [4] we have revised Ref. [3] and showed that α_s in the Bethe-Salpeter equations, including DGLAP, is basically replaced by the effective coupling α_s^{eff} given by much more complicated expression than Eq. (1):

$$\alpha_s^{eff} = \alpha_s(\mu^2) + \frac{1}{\pi b} \arctan \left(\frac{\pi [\ln(k_\perp^2/\beta\Lambda^2) - \ln(\mu^2/\Lambda^2)]}{\pi^2 + \ln(k_\perp^2/\beta\Lambda^2) \ln(\mu^2/\Lambda^2)} \right) \quad (2)$$

where the Sudakov variable β is the fraction of the longitudinal of the ladder parton, $\Lambda = \Lambda_{QCD}$ and $b = [11N - 2n_f]/(12\pi)$. However when μ , the starting point of the Q^2 -evolution is chosen large enough, namely when

$$\mu \gg \Lambda_{QCD} e^{\pi/2}, \quad (3)$$

Eq. (2) can be simplified down to

$$\alpha_s^{eff} \approx \alpha_s(k_\perp^2/\beta). \quad (4)$$

When Eq. (2) is applied to the DIS structure functions at $x \sim 1$ and Eq. (3) is also fulfilled, Eqs. (2,4) can be approximated by the DGLAP expression of Eq. (1).

II. PARAMETRIZATION OF α_s FOR QCD PROCESSES IN THE HARD KINEMATICS

Let us consider the contribution M_t of the Feynman graph depicted in Fig. 1. The cases with u and s -channel gluons factorized can be considered quite similarly. The solid lines in Fig. 1 denote quarks, though the generalization to the case of gluons is obvious. Through the paper we will assume that the lower particles, with momenta p_1, p'_1 , have small virtualities $\sim \mu^2$ whereas virtualities of the upper partons, with momenta q, q' are large: $-q^2 \sim -q'^2 \sim Q^2 \gg \mu^2$. Applying the Feynman rules to Fig. 1 we obtain:

$$M_t = i \frac{s^2}{4\pi^2} \int d\alpha d\beta dk_\perp^2 \frac{M(s, Q^2, s\alpha, s\beta, k_\perp^2)}{[s\beta - Q^2 - s\alpha\beta - k_\perp^2 + i\epsilon][s\alpha - s\alpha\beta - k_\perp^2 + i\epsilon]} \frac{\alpha_s(-s\alpha\beta - k_\perp^2)}{(s\alpha\beta + k_\perp^2 - i\epsilon)}. \quad (5)$$




FIG. 1: The right-hand side of Eq. (5)

We have used the standard Sudakov variables $k = -\alpha(q + xp) + \beta p + k_\perp$ and we have dropped the color factors as unessential for our analysis. M corresponds to the blob in Fig. 1. In Eq. (5) we have also neglected the virtuality $p^2 = \mu^2$ of the initial parton and denoted $s = 2pq$, $x = Q^2/2pq$, $Q^2 = -q^2$. Amplitude M is unknown, so it is impossible to perform the integration over any of the variables in Eq. (5). However, if we assume the leading logarithmic (LL) accuracy, we can use the QCD -generalization of the bremsstrahlung Gribov theorem. According to it, M does not depend on α and β . Integrating over α in Eq. (5) is conventionally performed with closing the integration contour down and taking the residue at $s\alpha = (+k_\perp^2 - i\epsilon)/(1 - \beta)$. It converts Eq. (5) into

$$M_t = -\frac{1}{2\pi} \int_{\mu^2}^s \frac{dk_\perp^2}{k_\perp^2} \int_{\beta_0}^1 d\beta \frac{(1-\beta)}{\beta} M(s, Q^2, k_\perp^2) \alpha_s \left(-\frac{k_\perp^2}{(1-\beta)} \right) \quad (6)$$

where $\beta_0 = x + k_\perp^2/s$. Obviously, $\beta_0 \approx x = Q^2/2pq$ when $x \sim 1$ and the upper limit s of the integration over $k_\perp^2$ can be changed for Q^2 . The minus sign of the α_s -argument in Eq. (6) indicates explicitly that the argument is space-like and for the space-like argument α_s is given by the well-known expression:

$$\alpha_s \left(-\frac{k_\perp^2}{(1-\beta)} \right) = \frac{1}{b \ln \left(k_\perp^2 / ((1-\beta)\Lambda^2) \right)}. \quad (7)$$

With the LL accuracy, $k_\perp^2/(1-\beta) \approx k_\perp^2$ and therefore in Eq. (6) $\alpha_s \approx \alpha_s(k_\perp^2)$. The minus sign of the argument of α_s is traditionally dropped, which drives us back to the standard expression of Eq. (1).

III. PARAMETRIZATION OF α_s IN THE BETHE-SALPETER EQUATIONS

In this section we study the parametrization of α_s in the Bethe-Salpeter equation for the forward scattering amplitude A . Let us assume that A obeys the following Bethe-Salpeter equation:

$$A = A_0 + \frac{i}{4\pi^2} \int dk_\perp^2 d\beta dm^2 M(s\beta, Q^2, (m^2\beta + k_\perp^2)) \frac{(1-\beta)k_\perp^2}{(m^2\beta + k_\perp^2 - i\epsilon)^2} \frac{\alpha_s(m^2)}{[m^2 + i\epsilon]}. \quad (8)$$

The second term in the rhs of Eq. (8) is depicted in Fig. 2. We have used the standard Sudakov variables: $k = -\alpha(q + xp) + \beta p + k_\perp$. Following Ref. [3], we have replaced the Sudakov variable $\alpha = 2pk/2pq$ by the new variable $m^2 = (p-k)^2$. M in Eq. (8) denotes the upper blob in Fig. 2. It includes both the off-shell amplitude A and a kernel. Now we just notice that Eq. (8) can be solved only after M has been known. A_0 stands for an inhomogeneous term. We focus on integrating over α in Eq. (8) and introduce

$$I = \int_{-\infty}^{\infty} dm^2 M(s\beta, Q^2, (m^2\beta + k_\perp^2)) \frac{(1-\beta)k_\perp^2}{(m^2\beta + k_\perp^2 - i\epsilon)^2} \frac{\alpha_s(m^2)}{[m^2 + i\epsilon]}. \quad (9)$$




FIG. 2: The integral contribution in Eq. (8)

The integrand of Eq. (9) has the singularities in m^2 . First, there are two poles from the propagators:

$$m^2 = -k_{\perp}^2/\beta + i\epsilon \quad (10)$$

and

$$m^2 = 0 - i\epsilon. \quad (11)$$


Second, there are two cuts. The first cut is originated by the k^2 -dependence of M . In particular, it can be the logarithmic dependence. The cut begins at

$$m^2 = -k_{\perp}^2/\beta + i\epsilon \quad (12)$$

and goes to the left. The second cut is related to α_s . It begins at

$$m^2 = 0 - i\epsilon \quad (13)$$

and goes to the right. The singularities (10-13) are depicted in Fig. 3. The integration over m^2 in Eq. (9) runs along

FIG. 3: Singularities of I given by Eqs. (10 - 13)

the $\Re m^2$ -axis from $-\infty$ to ∞ , so the integral can be calculated with choosing an appropriate closed integration contour C and taking residues. The contour C should include the line $-\infty < m^2 < \infty$ and a semi-circle C_R with radius R . The contour C_R may be situated either in the upper or in the right semi-plane of the m^2 -plane. However, if we choose C_R to be in the upper semi-plane, we should deal with the cut (12) of an unknown amplitude M , which is impossible without making assumptions about M . Alternatively, choosing the contour C_R in the lower semi-plane

involves analysis of the cut (13) of α_s and α_s is known. By this reason, we choose the latter option for C_R . the contour C_{cut} which runs along both sides of the cut (13). According to the Cauchy theorem,

$$I_C \equiv \int_C dm^2 M(s\beta, Q^2, k_\perp^2) \frac{(1-\beta)k_\perp^2}{(m^2\beta + k_\perp^2 - i\epsilon)^2} \frac{\alpha_s(m^2)}{[m^2 + i\epsilon]} = -2\pi i \frac{(1-\beta)}{k_\perp^2} M(s\beta, Q^2, -k_\perp^2/(1-\beta)) \alpha_s(\mu^2). \quad (14)$$

The rhs of Eq. (14) is the residue at the pole (11) and μ is introduced to regulate the IR singularity for α_s . It should be chosen as large as $\mu \gg \Lambda$ to guarantee applicability of the perturbative expression for α_s . When the initial partons are quarks, μ should also obey $\mu \gg$ the quark mass. Obviously,

$$I_C = I + I_{cut} + I_R \quad (15)$$

where I is defined in Eq. (9), I_R stands for the integration over the lower semi-circle and I_{cut} refers to the integration along the cut (13). I_R can be dropped because $I_R \rightarrow 0$ when $R \rightarrow \infty$. Now we specify I_{cut} :

$$I_{cut} = -2i \int_{\mu^2}^{\infty} dm^2 M(s\beta, Q^2, (m^2\beta + k_\perp^2)) \frac{(1-\beta)k_\perp^2}{(m^2\beta + k_\perp^2 - i\epsilon)^2} \frac{\Im \alpha_s(m^2)}{m^2}. \quad (16)$$

The integration in Eq. (16) cannot be done precisely because it involves the unknown amplitude M depending on m^2 . Nevertheless, it is possible to estimate I_{cut} . Indeed, the m^2 -dependence of M in Eq. (16) can be neglected in the region $m^2 \ll k_\perp^2/\beta$. Doing so, we obtain the following estimate of I :

$$I \approx -2i \frac{(1-\beta)}{k_\perp^2} M(s\beta, Q^2, k_\perp^2) \int_{\mu^2}^{k_\perp^2/\beta} dm^2 \frac{\Im \alpha_s(m^2)}{m^2} = -\frac{2i\pi(1-\beta)}{k_\perp^2} M(s\beta, Q^2, k_\perp^2) \alpha_s^{eff}, \quad (17)$$

with α_s^{eff} given by Eq. (4). When μ is chosen as large that Eq. (3) is fulfilled, we can drop π^2 in Eq. (17) and arrive at the estimate

$$I \approx -\frac{2i\pi(1-\beta)}{k_\perp^2} M(s\beta, Q^2, k_\perp^2) \alpha_s(k_\perp^2/\beta). \quad (18)$$

Finally, we give several estimates of $R = |\alpha_s^{eff} - \alpha_s|/\alpha_s^{eff}$, assuming that $\Lambda \approx 0.1\text{GeV}$: $R = 5\%$ at $\mu^2 \approx 30\text{GeV}^2$; then $R = 10\%$ at $\mu^2 \approx 2.4\text{GeV}^2$ and $R = 50\%$ at $\mu^2 \approx 0.8\text{GeV}^2$.

IV. ACKNOWLEDGMENTS

The work is partly supported by the EU grant MTKD-CT-2004-510126 in partnership with the CERN Physics Department and Russian State Grant for Scientific School RSGSS-5788.2006.2.

-
- [1] Yu.L. Dokshitzer, D.I. Diakonov, S.I. Troyan. Phys.Rep.58(1980)269.
 - [2] D. Amati, A. Bassetto, M. Ciafaloni, G. Marchesini, G. Veneziano. Nucl.Phys.B 173 (1980) 429; A. Bassetto, M. Ciafaloni, G. Marchesini. Phys. Rep. C 100 (1983) 201.
 - [3] Yu.L. Dokshitzer, D.V. Shirkov. Z. Phys. C 67 (1995) 449.
 - [4] B.I. Ermolaev, S.I. Troyan. Phys.Lett. B 666 (2008) 256.